

Brand Name: ACOFEN-P

Composition: Aceclofenac & Paracetamol IP. 325 mg tablet

Dosage Form: Tablets

Category: Non-Steroidal Anti-Inflammatory Drug (NSAID) with Analgesic

Packaging Type: Blister

Packaging: 20 x 10

Brief Product Introduction:

ACOFEN-P is a fast-acting and well-tolerated combination of **Aceclofenac and Paracetamol**, developed by **AstraEureka Pharmaceutical**, a trusted **manufacturer of Aceclofenac and Paracetamol composition**. This dual-action analgesic and anti-inflammatory formulation is widely used for the effective management of pain and inflammation in musculoskeletal and joint disorders. Aceclofenac provides targeted anti-inflammatory activity by inhibiting prostaglandin synthesis, while Paracetamol enhances the analgesic and antipyretic effect. ACOFEN-P is ideal for conditions such as arthritis, back pain, toothache, muscle sprain, and postoperative pain. Its balanced combination ensures rapid pain relief, improved joint mobility, and reduced inflammation without significant gastric irritation. Suitable for short-term and chronic use under medical supervision, ACOFEN-P is a preferred choice in orthopedic, dental, and general practice for daily pain management needs.

Composition:

Each uncoated tablet contains:

Aceclofenac IP.....100 mg

Paracetamol IP.....325 mg

Excipients.....q.s.

Key Benefits:

1. Effective pain relief with anti-inflammatory action
2. Reduces fever and inflammation simultaneously
3. Fast onset of action with extended symptom control
4. Lower gastric irritation compared to conventional NSAIDs
5. Well-suited for musculoskeletal and general pain conditions

Indications:

1. Rheumatoid arthritis, osteoarthritis, ankylosing spondylitis
2. Low back pain, cervical pain, muscle stiffness
3. Dental pain, post-extraction discomfort
4. Fever with body aches
5. Post-traumatic and postoperative pain

Directions for Use:

1. To be taken orally with water
2. Best taken after meals to avoid stomach upset
3. Swallow whole without breaking or crushing
4. Use only as per the dosage prescribed by a physician

Dosage and Administration:

1. **Adults:**
 - One tablet twice daily or as directed by the physician
 - Adjust dose based on clinical condition and response
 - Not suitable for pediatric use unless prescribed by a doctor

Safety Information:

1. To be used only under medical supervision
2. Avoid alcohol and NSAIDs during the course of therapy
3. Monitor liver function in prolonged use
4. Caution advised in patients with liver, kidney, or gastrointestinal conditions

Mechanism of Action:

- **Aceclofenac** inhibits the cyclooxygenase (COX) enzyme responsible for prostaglandin production, reducing pain and inflammation at the site of injury or inflammation.
- **Paracetamol** works centrally in the brain to reduce fever and increase the pain threshold, enhancing the overall analgesic effect.

Contraindications:

1. Known allergy to Aceclofenac, Paracetamol, or any excipients
2. Active peptic ulcer or gastrointestinal bleeding
3. Severe hepatic or renal impairment
4. Pregnancy and breastfeeding (unless recommended by a physician)

Side Effects:

1. Nausea, indigestion, abdominal pain
2. Drowsiness, dizziness
3. Rare: Rash, allergic reaction, liver enzyme elevation
4. Long-term use may lead to gastric or liver complications

Important Notice:

- Not for self-medication or prolonged unsupervised use
- Discontinue immediately in case of allergic symptoms
- Do not exceed the recommended dose
- Suitable lifestyle and physical therapy may improve treatment outcomes

Storage Condition:

1. Store below 25°C in a dry, cool place
2. Protect from moisture and direct sunlight
3. Keep out of reach of children

Packing Information:

- Each blister strip contains 10 uncoated tablets
- Packed in 20 x 10 tablets per box (blister pack)

