

Brand Name: NIMOON-P

Composition: Nimesulide BP 100 mg & Paracetamol IP 325 mg

Dosage Form: Tablets

Category: Non-Steroidal Anti-Inflammatory Drug (NSAID) / Analgesic-Antipyretic Combination

Packaging Type: BLISTER PACK

Packaging: 20 x 10 tablets

Brief Product Introduction:

NIMOON-P, manufactured by **AstraEureka Pharmaceutical**, a leading manufacturer of **Nimesulide and Paracetamol combination tablets**, is a dual-action analgesic formulated to provide effective relief from pain and fever. The tablet combines **Nimesulide 100 mg**, a selective COX-2 inhibitor with anti-inflammatory properties, and **Paracetamol 325 mg**, a well-established analgesic and antipyretic. NIMOON-P is indicated in conditions like musculoskeletal pain, dental pain, headache, backache, joint pain, fever, and inflammation. It offers rapid and long-lasting relief by acting at both central and peripheral pain pathways. The uncoated tablet ensures faster dissolution and absorption. It is a preferred choice for clinicians aiming for enhanced patient comfort in acute pain and febrile conditions.

Composition:

Each uncoated tablet contains:

Nimesulide BP.....	100 mg
Paracetamol IP.....	325 mg
Excipients.....	q.s.

Key Ingredients:

- Nimesulide BP 100 mg
- Paracetamol IP 325 mg

Key Benefits:

1. Effective relief from mild to moderate pain
2. Reduces fever quickly and efficiently
3. Combines anti-inflammatory and antipyretic action
4. Faster onset of action due to synergistic formulation
5. Suitable for musculoskeletal and dental indications

Indications:

1. Fever associated with body pain
2. Headache and migraine

3. Dental pain and inflammation
4. Musculoskeletal pain, sprain, and strain
5. Osteoarthritis and rheumatoid conditions
6. Dysmenorrhea (menstrual cramps)

Directions for Use:

1. Take orally with water after meals
2. Do not chew or crush
3. Follow physician's instructions strictly

Dosage and Administration:

1. **Adults:** One tablet twice daily or as prescribed
2. **Children:** Not recommended without physician supervision
3. Do not exceed 2 tablets in 24 hours without medical advice

Mechanism of Action:

- **Nimesulide** selectively inhibits the COX-2 enzyme, thereby reducing the synthesis of prostaglandins that cause pain and inflammation.
- **Paracetamol** acts centrally in the hypothalamus to reduce fever and inhibit pain transmission.
- Together, they deliver dual action—reducing both inflammation and central fever response—offering comprehensive symptomatic relief.

Safety Information:

1. Use under medical supervision only
2. Avoid use in liver impairment and gastrointestinal conditions
3. Not recommended for long-term therapy
4. Do not use with alcohol or hepatotoxic drugs

Contraindications:

1. Known hypersensitivity to Nimesulide, Paracetamol, or NSAIDs
2. Liver dysfunction or active liver disease
3. Peptic ulcers or gastrointestinal bleeding
4. Severe renal impairment
5. Children under 12 years (unless prescribed)

Side Effects:

1. Gastric irritation, nausea

2. Dizziness or headache
3. Elevated liver enzymes (with prolonged use)
4. Rare: skin rash, allergic reaction

Important Notice:

- For short-term use only
- Discontinue use if symptoms persist or worsen
- Avoid taking with other paracetamol-containing products
- Regular liver function monitoring recommended if used long term

Storage Condition:

1. Store in a cool, dry place below 25°C
2. Protect from light and moisture
3. Keep out of reach of children

Packing Information:

- Packed in blister strips
- Each pack contains 20 x 10 tablets (200 tablets per box)

