**Brand Name: PITOMOL 250 SUSPENSION** 

**Composition:** Paracetamol IP 250 mg

**Dosage Form: SUSPENSION Category:** Analgesic / Antipyretic

Packaging Type: PET BOTTLE WITH OUTER BOX

Packaging: 60 ML

#### **Brief Product Information:**

PITOMOL 250 SUSPENSION is a pediatric-friendly paracetamol formulation manufactured by AstraEureka Pharmaceutical, a trusted manufacturer of composition including Paracetamol IP. It is used for effective relief from fever and mild to moderate pain in children. This pleasant-flavored suspension ensures accurate tients. dosing and better compliance in pediatric patients.

### **Composition:**

#### Each 5 ml contains:

- Paracetamol IP......250 mg
- In a flavoured syrup base.....q.s.

# **Key Ingredients:**

- 1. Paracetamol IP 250 mg (per 5 ml): A widely used analgesic and antipyretic
- 2. Flavoured Syrup Base Q.S.: Palatable formulation for easy administration in children.

#### **Key Benefits:**

- 1. Reduces fever quickly and effectively.
- 2. Provides relief from mild to moderate pain including headache, toothache, sore throat, and post-vaccination discomfort.
- 3. Well-tolerated and safe when used as directed.
- 4. Sugar-based, child-friendly taste enhances acceptability.
- 5. Suitable for pediatric use under supervision.
- 6. Non-irritating to the gastric mucosa.

#### **Indications:**

- 1. Fever (any origin)
- 2. Mild to moderate pain in children
- 3. Teething-related discomfort
- 4. Headache or earache

- 5. Post-immunization fever or body ache
- 6. Sore throat or cold-associated fever

#### **Directions for Use:**

- 1. Shake the bottle well before use.
- 2. Use the calibrated measuring spoon or dropper provided.
- 3. Administer orally, preferably after food.
- 4. Follow dosing instructions strictly as per physician's guidance.
- 5. Do not exceed recommended frequency.

#### **Doses and Administration:**

- 1. Children (based on age and body weight):
  - 10–15 mg/kg body weight per dose every 4 to 6 hours.
  - Do not administer more than 4 doses in 24 hours.
- 2. **Infants and toddlers:** Only under medical supervision.
- 3. Always use the correct measuring device for accurate dosing.
- 4. Avoid prolonged or continuous use without physician advice.

### **Safety Information:**

- 1. Do not exceed recommended dose to avoid risk of liver damage.
- 2. Ensure proper dosing intervals (minimum 4 hours between doses).
- 3. Not for use in children under 3 months unless advised by a doctor.
- 4. Keep out of reach of children when not in use.
- 5. Avoid combining with other medications containing paracetamol.
- 6. Suitable for short-term use only unless directed otherwise.

#### **Mechanism of Action:**

- Paracetamol (acetaminophen) acts centrally in the brain to inhibit the enzyme cyclooxygenase (COX) involved in the synthesis of prostaglandins, which are responsible for pain and fever.
- It primarily acts on the **hypothalamic heat-regulating center** to reduce elevated body temperature.
- Unlike NSAIDs, it has minimal anti-inflammatory effects and is gentle on the stomach, making it ideal for children.

#### **Contraindications:**

- 1. Known hypersensitivity to paracetamol or any excipients.
- 2. Severe liver impairment or active liver disease.
- 3. Known hereditary fructose intolerance (due to sorbitol presence).

4. Use with caution in patients with renal impairment.

## **Side Effects:**

- 1. Rare: Allergic reactions (rash, itching, swelling).
- 2. Liver toxicity in case of overdose or prolonged use.
- 3. Very rare: Blood disorders like thrombocytopenia.
- 4. Gastrointestinal discomfort (rare).

# **Important Notice:**

- Always check the label for correct dose before use.
- Do not use in combination with other paracetamol-containing products.
- In case of overdose, seek immediate medical help—even if no symptoms

PHARMACEU

- Stop use and consult a physician if fever or pain persists beyond 3 days.
- Store and administer with proper pediatric guidance.

### **Storage Condition:**

- 1. Store below 25°C in a cool and dry place.
- 2. Protect from light and moisture.
- 3. Do not freeze.
- 4. Keep the cap tightly closed after each use.
- 5. Keep out of reach of children.

# **Packing Information:**

- Packaging: 60 ML