#### **Brand Name: MEDPROSTO TABLET**

**Composition:** Medroxyprogesterone Acetate Tablet IP

**Dosage Form:** TABLET

**Category:** Hormonal Therapy / Progestin

Packaging Type: ALU-ALU **Packaging:** 10 x 10 TABLETS

#### **Brief Product Introduction:**

MEDPROSTO TABLET is a progestin-based oral formulation manufactured by AstraEureka Pharmaceutical, indicated for the management of various menstrual disorders, endometriosis, and hormone replacement therapy. Each tablet contains 10 mg of **Medroxyprogesterone Acetate**, a synthetic derivative of progesterone, used to regulate ovulation and menstrual cycles.

#### **Composition:**

#### **Each Uncoated Tablet Contains:**

regulate ovulation and menstrual cycles	
1311	Produce
Composition:	Todatels c
Each Uncoated Tablet Contains:	a The
Medroxyprogesterone Acetate IP	10 mg
Excipients	q.s.

#### **Key Ingredients:**

- Medroxyprogesterone Acetate IP (10 mg):
  - Synthetic progestin with properties similar to natural progesterone
  - Helps restore hormonal balance in women with menstrual irregularities

PHARMACEUTICALS

### **Key Benefits:**

- 1. Regulates irregular menstrual cycles
- 2. Treats secondary amenorrhea (absence of menstruation)
- 3. Effective in controlling abnormal uterine bleeding due to hormonal imbalance
- 4. Used in hormone replacement therapy (HRT) in postmenopausal women
- 5. Supports management of endometriosis and some hormone-responsive
- 6. Can be used as part of contraceptive regimens under medical guidance

#### **Indications:**

- 1. Secondary Amenorrhea
- 2. Abnormal Uterine Bleeding due to hormonal imbalance
- 3. Endometriosis
- 4. Hormone Replacement Therapy in postmenopausal women
- 5. As an adjunct in treatment of endometrial carcinoma or renal carcinoma (as prescribed)

6. Prevention of endometrial hyperplasia during estrogen therapy

#### **Directions for Use:**

- 1. To be taken orally with or without food
- 2. Follow the exact dosage schedule as prescribed by the physician
- 3. Swallow whole with water, do not crush or chew

#### **Dosage and Administration:**

- 1. **Secondary Amenorrhea:** 5–10 mg once daily for 5–10 days
- 2. **Abnormal Uterine Bleeding:** 5–10 mg daily for 5–10 days during the second half of the menstrual cycle
- 3. **Endometriosis:** 10 mg three times daily for 90 days beginning on the first day of the menstrual cycle
- 4. **Hormone Replacement Therapy (HRT):** Dose varies based on estrogen combination; typically 5–10 mg daily for 12–14 days per cycle
- 5. Dosage must be individualized as per patient condition and medical advice

### Safety Information:

- 1. To be used under medical supervision
- 2. Not recommended for long-term use without regular follow-up
- 3. Regular gynecological examinations advised during prolonged therapy
- 4. Use with caution in patients with a history of depression or thromboembolic disorders

#### **Mechanism of Action:**

- Medroxyprogesterone Acetate binds to progesterone receptors in the uterus and other target tissues.
- It induces secretory changes in the endometrium, reduces pituitary gonadotropin secretion, and inhibits ovulation when taken in sufficient doses.
- It also modulates the hormonal balance by opposing estrogen effects on the endometrial lining, thereby stabilizing the menstrual cycle.

#### **Contraindications:**

- 1. Known hypersensitivity to Medroxyprogesterone or any component of the tablet
- 2. Pregnancy (Category X contraindicated)
- 3. Undiagnosed vaginal bleeding
- 4. Active or history of thromboembolic disorders (DVT, PE)
- 5. Severe hepatic dysfunction or liver disease

6. Known or suspected breast or genital malignancy

#### **Side Effects:**

- 1. Nausea, bloating, weight gain
- 2. Breast tenderness
- 3. Headache, mood swings, depression
- 4. Irregular spotting or breakthrough bleeding
- 5. Fatigue, dizziness
- 6. Rare: thromboembolic events, allergic reactions

### **Important Notice:**

- Hormonal therapy should only be initiated after a thorough clinical evaluation
- Not intended for use as a routine contraceptive unless specifically prescribed
- Regular monitoring is advised during long-term therapy

## **Storage Condition:**

- 1. Store in a cool and dry place, below 25°C
- 2. Protect from light and moisture
- 3. Keep out of reach of children

# **Packing Information:**

- Supplied in ALU-ALU packs of 10 x 10 tablets A C E U
- Clearly labeled with batch number, manufacturing and expiry date

Tusted Quality Products From P