

**Brand Name: ROJASTA-20 GOLD CAPSULE**

**Composition:** Rosuvastatin 20 mg, Aspirin IP 75 mg & Clopidogrel 75 mg

**Dosage Form:** Capsule

**Category:** Cardiovascular – Antihyperlipidemic & Antiplatelet Combination

**Packaging Type:** Alu-Alu

**Packaging:** 10 capsules per strip × 10 strips per box (10 x 10)

**Brief Product Introduction:**

**ROJASTA-20 GOLD** is a triple-combination cardiovascular capsule manufactured by **AstraEureka Pharmaceutical**, a trusted **manufacturer of composition** for cardiometabolic therapy. This fixed-dose combination includes **Rosuvastatin (20 mg)**, a potent HMG-CoA reductase inhibitor that lowers LDL cholesterol and triglycerides; **Aspirin (75 mg)**, an antiplatelet agent that prevents clot formation; and **Clopidogrel (75 mg)**, a P2Y<sub>12</sub> receptor antagonist that further inhibits platelet aggregation. ROJASTA-20 GOLD is indicated for the prevention of heart attacks, strokes, and complications in patients with established cardiovascular disease, diabetes, or hyperlipidemia. The once-daily capsule simplifies complex regimens and improves patient compliance. It offers synergistic protection by managing dyslipidemia while also reducing thrombotic risk, making it essential in secondary prevention protocols and post-angioplasty care.

**Composition:**

**Each hard gelatin capsule contains:**

Rosuvastatin Calcium IP

equivalent to Rosuvastatin.....20 mg

Aspirin IP.....75 mg

Clopidogrel Bisulphate IP

equivalent to Clopidogrel.....75 mg

Excipients.....q.s.

**Key Ingredients:**

- **Rosuvastatin (20 mg):** Lowers LDL, total cholesterol, and triglycerides
- **Aspirin (75 mg):** Inhibits thromboxane A<sub>2</sub>, reducing platelet aggregation
- **Clopidogrel (75 mg):** Blocks ADP receptors on platelets, preventing clot formation

**Key Benefits:**

1. Comprehensive cardiovascular protection
2. Reduces risk of heart attack, stroke, and stent thrombosis
3. Controls cholesterol and prevents atherosclerotic progression

4. Minimizes platelet aggregation with dual antiplatelet action
5. Once-daily dosing improves treatment adherence
6. Essential in post-angioplasty and post-MI maintenance therapy

### **Indications:**

1. Secondary prevention of cardiovascular events (MI, stroke)
2. Patients with coronary artery disease, peripheral artery disease, or recent angioplasty
3. High-risk dyslipidemic patients requiring antiplatelet therapy
4. Atherosclerosis prevention and plaque stabilization

### **Directions for Use:**

1. Take one capsule daily with or after meals
2. Swallow whole with water; do not crush or chew
3. Follow the dosage exactly as prescribed by the physician

### **Dosage and Administration:**

1. **Adults:** One capsule once daily or as directed by the healthcare provider
2. Dose adjustments may be based on lipid levels, platelet function, and tolerance
3. Regular lipid profile and platelet monitoring are recommended

### **Safety Information:**

1. Use with caution in patients with bleeding disorders or liver impairment
2. Monitor for signs of GI bleeding or muscle toxicity
3. Avoid alcohol and NSAIDs during treatment
4. Not recommended during pregnancy or breastfeeding unless advised by a physician
5. Do not discontinue abruptly without medical supervision

### **Mechanism of Action:**

- **Rosuvastatin** inhibits HMG-CoA reductase, reducing cholesterol synthesis in the liver and increasing LDL receptor uptake
- **Aspirin** irreversibly inhibits COX-1 in platelets, blocking thromboxane A<sub>2</sub>, and reducing platelet aggregation
- **Clopidogrel** inhibits ADP binding to P2Y<sub>12</sub> receptors on platelets, preventing activation and cross-linking, thus reducing clot formation
- Together, they provide lipid control, anti-inflammatory effects, and antithrombotic protection

**Contraindications:**

1. Known hypersensitivity to Rosuvastatin, Aspirin, Clopidogrel, or any excipient
2. Active peptic ulcer or gastrointestinal bleeding
3. Hemorrhagic stroke or bleeding disorders
4. Severe hepatic or renal impairment
5. Pregnancy and lactation

**Side Effects:**

1. Myalgia, headache, dizziness
2. Gastric irritation, nausea, dyspepsia
3. Risk of GI bleeding or bruising
4. Rare: Rhabdomyolysis, liver enzyme elevation, hypersensitivity reactions

**Important Notice:**

- This is a **prescription-only medicine** and should be taken under medical supervision.
- Lifestyle modifications, including diet and exercise, are essential for optimal cardiovascular protection.
- Do not discontinue therapy without physician approval.

**Storage Condition:**

1. Store in a cool, dry place below 25°C
2. Protect from moisture and direct sunlight
3. Keep in original alu-alu pack until administration

**Packing Information:**

- **Packaging:** 10 capsules per alu-alu strip × 10 strips per carton (10 x 10)
- **Net Quantity:** 100 capsules per box